

References:

- Acton, T. B., Mead, J., Steiner, A. M., and Vershon, A. K. (2000). Scanning mutagenesis of Mcm1: residues required for DNA binding, DNA bending, and transcriptional activation by a MADS-box protein., *Mol Cell Biol* 20, 1-11.
- el-Deiry, W. S., Kern, S. E., Pietenpol, J. A., Kinzler, K. W., and Vogelstein, B. (1992). Definition of a consensus binding site for p53, *Nat Genet* 1, 45-49.
- Passmore, S., Elble, R., and Tye, B. K. (1989). A protein involved in minichromosome maintenance in yeast binds a transcriptional enhancer conserved in eukaryotes, *Genes Dev* 3, 921-935.
- Rusmann, C., Truss, M., Fix, A., Naumer, C., Herrmann, T., Schmitt, J., Stollhof, J., Beigang, R., and Beato, M. (1997). Crosslinking of progesterone receptor to DNA using tuneable nanosecond, picosecond and femtosecond UV laser pulses, *Nucleic Acids Res* 25, 2478-2484.
- Saito, I., and Matsuura, T. (1985). Chemical aspects of UV-induced cross-linking of proteins to nucleic acids. Photoreactions with lysine and tryptophan, *Acc Chem Res* 18, 134-141.
- Tan, S., and Richmond, T. J. (1998). Crystal structure of the yeast MAT α 2/MCM1/DNA ternary complex, *Nature* 39, 660-666.
- Wyrick, J. J., Aparicio, J. G., Chen, T., Barnett, J. D., Jennings, E. G., Young, R. A., Bell, S. P., and Aparicio, O. M. (2001). Genome-wide distribution of ORC and MCM proteins in *S. cerevisiae*: high-resolution mapping of replication origins, *Science* 294, 2357-2360.